

Machine Learning



Support Vector Machines

Eric Xing




Lecture 4, August 12, 2010

Reading:


© Eric Xing @ CMU, 2006-2010



What is a good Decision Boundary?




- Why we may have such boundaries?
 - Irregular distribution
 - Imbalanced training sizes
 - outliers




Classification and Margin

- Parameterizing decision boundary
 - Let w denote a vector orthogonal to the decision boundary, and b denote a scalar "offset" term, then we can write the decision boundary as:

$$w^T x + b = 0$$






Classification and Margin

- Parameterizing decision boundary

- Let w denote a vector orthogonal to the decision boundary, and b denote a scalar "offset" term, then we can write the decision boundary as:

$$\frac{w^T}{\|w^T\|} x + \frac{b}{\|w^T\|} = 0$$



- Margin

$$(w^T x_i + b) / \|w\| > +c / \|w\| \quad \text{for all } x_i \text{ in class 2}$$


$$(w^T x_i + b) / \|w\| < -c / \|w\| \quad \text{for all } x_i \text{ in class 1}$$

Or more compactly:

$$(w^T x_i + b) y_i / \|w\| > c / \|w\|$$

The margin between two points

$$m = d^- + d^+ =$$



Maximum Margin Classification

- The margin is:

$$m = \frac{w^T}{\|w\|} (x_{i^*} - x_{j^*}) = \frac{2c}{\|w\|}$$

- Here is our Maximum Margin Classification problem:

$$\begin{aligned} \max_w & \frac{2c}{\|w\|} \\ \text{s.t} & y_i (w^T x_i + b) / \|w\| \geq c / \|w\|, \quad \forall i \end{aligned}$$

Maximum Margin Classification, con'd.



- The optimization problem:

$$\begin{aligned} \max_{w,b} \quad & \frac{c}{\|w\|} \\ \text{s.t} \quad & y_i(w^T x_i + b) / \|w\| \geq c / \|w\|, \quad \forall i \end{aligned}$$

- But note that the magnitude of c merely scales w and b , and does not change the classification boundary at all! (why?)
- So we instead work on this cleaner problem:

$$\begin{aligned} \max_{w,b} \quad & \frac{1}{\|w\|} \\ \text{s.t} \quad & y_i(w^T x_i + b) \geq 1, \quad \forall i \end{aligned}$$

- The solution to this leads to the famous **Support Vector Machines** -- believed by many to be the best "off-the-shelf" supervised learning algorithm



Digression to Lagrangian Duality

- The Primal Problem

Primal:

$$\begin{aligned} \min_w \quad & f(w) \\ \text{s.t.} \quad & g_i(w) \leq 0, \quad i = 1, \dots, k \\ & h_i(w) = 0, \quad i = 1, \dots, l \end{aligned}$$

The generalized Lagrangian:

$$\mathcal{L}(w, \alpha, \beta) = f(w) + \sum_{i=1}^k \alpha_i g_i(w) + \sum_{i=1}^l \beta_i h_i(w)$$


the α 's ($\alpha_i \geq 0$) and β 's are called the Lagrangian multipliers

Lemma:

$$\max_{\alpha, \beta, \alpha_i \geq 0} \mathcal{L}(w, \alpha, \beta) = \begin{cases} f(w) & \text{if } w \text{ satisfies primal constraint s} \\ \infty & \text{o/w} \end{cases}$$

A re-written Primal:

$$\min_w \max_{\alpha, \beta, \alpha_i \geq 0} \mathcal{L}(w, \alpha, \beta)$$



Lagrangian Duality, cont.

- Recall the Primal Problem:

$$\min_w \max_{\alpha, \beta, \alpha_i \geq 0} \mathcal{L}(w, \alpha, \beta)$$

- The Dual Problem:

$$\max_{\alpha, \beta, \alpha_i \geq 0} \min_w \mathcal{L}(w, \alpha, \beta)$$


- Theorem (weak duality):**

$$d^* = \max_{\alpha, \beta, \alpha_i \geq 0} \min_w \mathcal{L}(w, \alpha, \beta) \leq \min_w \max_{\alpha, \beta, \alpha_i \geq 0} \mathcal{L}(w, \alpha, \beta) = p^*$$

- Theorem (strong duality):**

Iff there exist a saddle point of $\mathcal{L}(w, \alpha, \beta)$, we have

$$d^* = p^*$$



The KKT conditions

- If there exists some saddle point of \mathcal{L} , then the saddle point satisfies the following "Karush-Kuhn-Tucker" (KKT) conditions:

$$\frac{\partial}{\partial w_i} \mathcal{L}(w, \alpha, \beta) = 0, \quad i = 1, \dots, k$$


$$\frac{\partial}{\partial \beta_i} \mathcal{L}(w, \alpha, \beta) = 0, \quad i = 1, \dots, l$$

$$\alpha_i g_i(w) = 0, \quad i = 1, \dots, m$$

$$g_i(w) \leq 0, \quad i = 1, \dots, m$$

$$\alpha_i \geq 0, \quad i = 1, \dots, m$$

- **Theorem:** If w^* , α^* and β^* satisfy the KKT condition, then it is also a solution to the primal and the dual problems.



Solving optimal margin classifier

- Recall our opt problem:

$$\begin{aligned} \max_{w,b} \quad & \frac{1}{\|w\|} \\ \text{s.t} \quad & y_i(w^T x_i + b) \geq 1, \quad \forall i \end{aligned}$$


- This is equivalent to

$$\begin{aligned} \min_{w,b} \quad & \frac{1}{2} w^T w \\ \text{s.t} \quad & 1 - y_i(w^T x_i + b) \leq 0, \quad \forall i \end{aligned} \quad (*)$$

- Write the Lagrangian:

$$\mathcal{L}(w, b, \alpha) = \frac{1}{2} w^T w - \sum_{i=1}^m \alpha_i [y_i(w^T x_i + b) - 1]$$

- Recall that (*) can be reformulated as $\min_{w,b} \max_{\alpha_i \geq 0} \mathcal{L}(w, b, \alpha)$
Now we solve its **dual problem**: $\max_{\alpha_i \geq 0} \min_{w,b} \mathcal{L}(w, b, \alpha)$



The Dual Problem

$$\mathcal{L}(w, b, \alpha) = \frac{1}{2} w^T w - \sum_{i=1}^m \alpha_i [y_i (w^T x_i + b) - 1]$$

$$\max_{\alpha_i \geq 0} \min_{w, b} \mathcal{L}(w, b, \alpha)$$

- We minimize \mathcal{L} with respect to w and b first:

$$\nabla_w \mathcal{L}(w, b, \alpha) = w - \sum_{i=1}^m \alpha_i y_i x_i = 0, \quad (*)$$


$$\nabla_b \mathcal{L}(w, b, \alpha) = \sum_{i=1}^m \alpha_i y_i = 0, \quad (**)$$

Note that (*) implies:

$$w = \sum_{i=1}^m \alpha_i y_i x_i \quad (***)$$

- Plug (***) back to \mathcal{L} , and using (**), we have:

$$\mathcal{L}(w, b, \alpha) = \sum_{i=1}^m \alpha_i - \frac{1}{2} \sum_{i,j=1}^m \alpha_i \alpha_j y_i y_j (\mathbf{x}_i^T \mathbf{x}_j)$$



The Dual problem, cont.

- Now we have the following dual opt problem:

$$\max_{\alpha} \mathcal{J}(\alpha) = \sum_{i=1}^m \alpha_i - \frac{1}{2} \sum_{i,j=1}^m \alpha_i \alpha_j y_i y_j (\mathbf{x}_i^T \mathbf{x}_j)$$

$$\text{s.t. } \alpha_i \geq 0, \quad i = 1, \dots, k$$


$$\sum_{i=1}^m \alpha_i y_i = 0.$$

- This is, (again,) a **quadratic programming** problem.

- A global maximum of α_i can always be found.
- But what's the big deal??
- Note two things:

1. w can be recovered by $w = \sum_{i=1}^m \alpha_i y_i \mathbf{x}_i$ See next ...


2. The "kernel" $\mathbf{x}_i^T \mathbf{x}_j$ More later ...




I. Support vectors

- Note the KKT condition --- only a few α_i 's can be nonzero!!

$$\alpha_i g_i(w) = 0, \quad i = 1, \dots, m$$



Call the training data points whose α_i 's are nonzero the **support vectors (SV)**



Support vector machines

- Once we have the Lagrange multipliers $\{\alpha_i\}$, we can reconstruct the parameter vector w as a weighted combination of the training examples:

$$w = \sum_{i \in SV} \alpha_i y_i \mathbf{x}_i$$

- For testing with a new data z

- Compute

$$w^T z + b = \sum_{i \in SV} \alpha_i y_i (\mathbf{x}_i^T z) + b$$

and classify z as class 1 if the sum is positive, and class 2 otherwise


- Note: w need not be formed explicitly

Interpretation of support vector machines




- The optimal \mathbf{w} is a linear combination of a small number of data points. This “sparse” representation can be viewed as data compression as in the construction of kNN classifier
- To compute the weights $\{\alpha_i\}$, and to use support vector machines we need to specify only the inner products (or kernel) between the examples $\mathbf{x}_i^T \mathbf{x}_j$
- We make decisions by comparing each new example \mathbf{z} with only the support vectors:

$$y^* = \text{sign} \left(\sum_{i \in SV} \alpha_i y_i (\mathbf{x}_i^T \mathbf{z}) + b \right)$$





II. The Kernel Trick

- Is this data linearly-separable?



- How about a quadratic mapping $\phi(x_i)$?





II. The Kernel Trick


- Recall the SVM optimization problem

$$\max_{\alpha} \quad \mathcal{J}(\alpha) = \sum_{i=1}^m \alpha_i - \frac{1}{2} \sum_{i,j=1}^m \alpha_i \alpha_j y_i y_j (\mathbf{x}_i^T \mathbf{x}_j)$$

$$\text{s.t.} \quad 0 \leq \alpha_i \leq C, \quad i = 1, \dots, m$$

$$\sum_{i=1}^m \alpha_i y_i = 0.$$

- The data points only appear as **inner product**
- As long as we can calculate the inner product in the feature space, we do not need the mapping explicitly
- Many common geometric operations (angles, distances) can be expressed by inner products
- Define the kernel function K by $K(\mathbf{x}_i, \mathbf{x}_j) = \phi(\mathbf{x}_i)^T \phi(\mathbf{x}_j)$



II. The Kernel Trick

- Computation depends on feature space
 - Bad if its dimension is much larger than input space

$$\max_{\alpha} \sum_{i=1}^m \alpha_i - \frac{1}{2} \sum_{i,j=1}^m \alpha_i \alpha_j y_i y_j K(\mathbf{x}_i, \mathbf{x}_j)$$


$$\text{s.t. } \alpha_i \geq 0, \quad i = 1, \dots, k$$

$$\sum_{i=1}^m \alpha_i y_i = 0.$$

Where $K(\mathbf{x}_i, \mathbf{x}_j) = \phi(\mathbf{x}_i)^t \phi(\mathbf{x}_j)$

$$y^*(z) = \text{sign} \left(\sum_{i \in SV} \alpha_i y_i K(\mathbf{x}_i, z) + b \right)$$

Transforming the Data



Note: feature space is of higher dimension than the input space in practice

- Computation in the feature space can be costly because it is high dimensional
 - The feature space is typically infinite-dimensional!
- The kernel trick comes to rescue

An Example for feature mapping and kernels



- Consider an input $\mathbf{x}=[x_1, x_2]$
- Suppose $\phi(\cdot)$ is given as follows

$$\phi\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = 1, \sqrt{2}x_1, \sqrt{2}x_2, x_1^2, x_2^2, \sqrt{2}x_1x_2$$

- An inner product in the feature space is

$$\left\langle \phi\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right), \phi\left(\begin{bmatrix} x_1' \\ x_2' \end{bmatrix}\right) \right\rangle =$$

- So, if we define the **kernel function** as follows, there is no need to carry out $\phi(\cdot)$ explicitly

$$K(\mathbf{x}, \mathbf{x}') = \left(1 + \mathbf{x}^T \mathbf{x}'\right)^2$$

More examples of kernel functions



- Linear kernel (we've seen it)

$$K(\mathbf{x}, \mathbf{x}') = \mathbf{x}^T \mathbf{x}'$$

- Polynomial kernel (we just saw an example)


$$K(\mathbf{x}, \mathbf{x}') = \left(\mathbf{1} + \mathbf{x}^T \mathbf{x}' \right)^p$$

where $p = 2, 3, \dots$. To get the feature vectors we concatenate all p th order polynomial terms of the components of \mathbf{x} (weighted appropriately)

- Radial basis kernel

$$K(\mathbf{x}, \mathbf{x}') = \exp\left(-\frac{1}{2} \|\mathbf{x} - \mathbf{x}'\|^2\right)$$

In this case the feature space consists of functions and results in a non-parametric classifier.



The essence of kernel

- Feature mapping, but “without paying a cost”

- E.g., polynomial kernel

$$K(x, z) = (x^T z + c)^d$$

- How many dimensions we’ve got in the new space?
- How many operations it takes to compute K()?

- Kernel design, any principle?

- K(x,z) can be thought of as a similarity function between x and z
- This intuition can be well reflected in the following “Gaussian” function (Similarly one can easily come up with other K() in the same spirit)

$$K(x, z) = \exp\left(-\frac{\|x - z\|^2}{2\sigma^2}\right)$$

- Is this necessarily lead to a “legal” kernel?



(in the above particular case, K() is a legal one, do you know how many dimension $\phi(x)$ is?)




Kernel matrix

- Suppose for now that K is indeed a valid kernel corresponding to some feature mapping ϕ , then for x_1, \dots, x_m , we can compute an $m \times m$ matrix $K = \{K_{i,j}\}$, where $K_{i,j} = \phi(x_i)^T \phi(x_j)$
- This is called a **kernel matrix!**
- Now, if a kernel function is indeed a valid kernel, and its elements are dot-product in the transformed feature space, it must satisfy:
 - Symmetry $K=K^T$
proof $K_{i,j} = \phi(x_i)^T \phi(x_j) = \phi(x_j)^T \phi(x_i) = K_{j,i}$
 - Positive –semidefinite $y^T K y \geq 0 \quad \forall y$
proof?
 - Mercer's theorem


SVM examples




linear



2^{nd} order polynomial





4^{th} order polynomial



8^{th} order polynomial


Examples for Non Linear SVMs – Gaussian Kernel






Example Kernel

- x_i is a bag of words
- Define $\phi(x_i)$ as a count of every n-gram up to $n=k$ in x_i .
 - This is huge space 26^k
 - What are we measuring by $\phi(x_i)^t \phi(x_j)$?
- Can we compute the same quantity on input space?
 - Efficient linear dynamic program!
- Kernel is a measure of similarity
- Must be positive semi-definite



Non-linearly Separable Problems



- We allow “error” ξ_i in classification; it is based on the output of the discriminant function $w^T x + b$
- ξ_i approximates the number of misclassified samples




Soft Margin Hyperplane

- Now we have a slightly different opt problem:

$$\begin{aligned} \min_{w,b} \quad & \frac{1}{2} w^T w + C \sum_{i=1}^m \xi_i \\ \text{s.t.} \quad & y_i (w^T x_i + b) \geq 1 - \xi_i, \quad \forall i \\ & \xi_i \geq 0, \quad \forall i \end{aligned}$$

- ξ_i are “slack variables” in optimization
- Note that $\xi_i=0$ if there is no error for \mathbf{x}_i
- ξ_i is an upper bound of the number of errors
- C : tradeoff parameter between error and margin

Hinge Loss





- Remember Ridge regression
 - Min [squared loss + $\lambda w^t w$]
- How about SVM?

$$\text{argmin}_{\{w,b\}} \underbrace{w^t w}_{\text{regularization}} + \lambda \underbrace{\sum_1^m \max(1 - y_i(w^t x_i + b), 0)}_{\text{Loss: hinge loss}}$$

regularization

Loss: hinge loss





The Optimization Problem


- The dual of this new constrained optimization problem is

$$\max_{\alpha} \quad \mathcal{J}(\alpha) = \sum_{i=1}^m \alpha_i - \frac{1}{2} \sum_{i,j=1}^m \alpha_i \alpha_j y_i y_j (\mathbf{x}_i^T \mathbf{x}_j)$$

$$\text{s.t.} \quad 0 \leq \alpha_i \leq C, \quad i = 1, \dots, m$$

$$\sum_{i=1}^m \alpha_i y_i = 0.$$

- This is very similar to the optimization problem in the linear separable case, except that there is an upper bound C on α_i now
- Once again, a QP solver can be used to find α_i





The SMO algorithm


- Consider solving the **unconstrained** opt problem:

$$\max_{\alpha} W(\alpha_1, \alpha_2, \dots, \alpha_m)$$

- We've already seen several opt algorithms!
 - ?
 - ?
 - ?
- Coordinate ascend:

Coordinate ascend





Sequential minimal optimization

- Constrained optimization:


$$\max_{\alpha} \quad \mathcal{J}(\alpha) = \sum_{i=1}^m \alpha_i - \frac{1}{2} \sum_{i,j=1}^m \alpha_i \alpha_j y_i y_j (\mathbf{x}_i^T \mathbf{x}_j)$$

$$\text{s.t.} \quad 0 \leq \alpha_i \leq C, \quad i = 1, \dots, m$$

$$\sum_{i=1}^m \alpha_i y_i = 0.$$

- Question: can we do coordinate along one direction at a time (i.e., hold all $\alpha_{[-i]}$ fixed, and update α_i ?)


The SMO algorithm



Repeat till convergence

1. Select some pair α_i and α_j to update next (using a heuristic that tries to pick the two that will allow us to make the biggest progress towards the global maximum).
2. Re-optimize $J(\alpha)$ with respect to α_i and α_j , while holding all the other α_k 's ($k \neq i; j$) fixed.

Will this procedure converge?



Convergence of SMO


$$\max_{\alpha} \mathcal{J}(\alpha) = \sum_{i=1}^m \alpha_i - \frac{1}{2} \sum_{i,j=1}^m \alpha_i \alpha_j y_i y_j (\mathbf{x}_i^T \mathbf{x}_j)$$

KKT:

$$\text{s.t. } 0 \leq \alpha_i \leq C, \quad i = 1, \dots, k$$

$$\sum_{i=1}^m \alpha_i y_i = 0.$$

- Let's hold $\alpha_3, \dots, \alpha_m$ fixed and reopt \mathcal{J} w.r.t. α_1 and α_2



Convergence of SMO

- The constraints:

$$\alpha_1 y_1 + \alpha_2 y_2 = \xi$$


$$0 \leq \alpha_1 \leq C$$


$$0 \leq \alpha_2 \leq C$$

- The objective:

$$\mathcal{J}(\alpha_1, \alpha_2, \dots, \alpha_m) = \mathcal{J}((\xi - \alpha_2 y_2) y_1, \alpha_2, \dots, \alpha_m)$$

- Constrained opt:






Cross-validation error of SVM

- The leave-one-out cross-validation error does not depend on the dimensionality of the feature space but only on the # of support vectors!

$$\text{Leave - one - out CV error} = \frac{\# \text{ support ve ctors}}{\# \text{ of training examples}}$$



Summary



- Max-margin decision boundary
- Constrained convex optimization
 - Duality
 - The KKT conditions and the support vectors
 - Non-separable case and slack variables
 - The SMO algorithm